

NEW UPPER AND LOWER BOUNDS FOR THE TRAPEZOID INEQUALITY OF ABSOLUTELY CONTINUOUS FUNCTIONS AND APPLICATIONS

MOHAMMAD W. ALOMARI

ABSTRACT. In this paper, new upper and lower bounds for the Trapezoid inequality of absolutely continuous functions are obtained. Applications to some special means are provided as well.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$, with $a < b$. The following inequality, known as *Hermite–Hadamard inequality* for convex functions, holds:

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}.$$

On the other hand, a very related inequality to (1.1) was known in literature as the ‘*Trapezoid inequality*’, which states that: if $f : [a, b] \rightarrow \mathbb{R}$ is twice differentiable such that $\|f''\|_\infty := \sup_{t \in (a, b)} |f''(t)| < \infty$, then

$$(1.2) \quad \left| \int_a^b f(x) dx - (b-a) \frac{f(a) + f(b)}{2} \right| \leq \frac{(b-a)^3}{12} \|f''\|_\infty.$$

In recent years many authors have established several inequalities connected to the Hermite–Hadamard’s inequality. For recent results, refinements, counterparts, generalizations and new Hermite–Hadamard-type inequalities under various assumptions for the functions involved the reader may refer to [1] – [17] and the references therein.

In this paper, new upper and lower bounds for the Trapezoid inequality of absolutely continuous functions are established. Applications to some special means are provided as well.

2. THE RESULT

Theorem 1. *Let $f : I \subset \mathbb{R} \rightarrow \mathbb{R}_+$ be an absolutely continuous mapping on I° , the interior of the interval I , where $a, b \in I$ with $a < b$. Then there exists $x \in (a, b)$*

Date: November 6, 2014.

2010 Mathematics Subject Classification. 26D15, 26D10.

Key words and phrases. Hermite–Hadamard inequality, Trapezoid inequality.

such that the double inequality

$$\begin{aligned}
 (2.1) \quad & \frac{(b-a)^2}{2M^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{M^2}{(x-a)(b-x)} f(x) \right] \\
 & \leq \frac{f(b) + f(a)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \\
 & \leq \frac{(b-a)^2}{2m^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{m^2}{(x-a)(b-x)} f(x) \right]
 \end{aligned}$$

holds, where

$$M := \left[\frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right],$$

and

$$m := \left[\frac{b-a}{2} - \left| x - \frac{a+b}{2} \right| \right].$$

Proof. Consider the function $F : [a, b] \rightarrow (0, \infty)$ defined by

$$F(t) = \frac{1}{b-t} \int_t^b f(s) ds - \frac{1}{t-a} \int_a^t f(s) ds$$

for all $t \in [a, b]$. Since f is absolutely continuous on $[a, b]$, then it is easy to see that F is differentiable on (a, b) . So that by the Mean-Value Theorem there exists $x \in (a, b)$ such that

$$(2.2) \quad F'(x) = \frac{F(b) - F(a)}{b-a}.$$

Simple calculations yield that

$$F'(x) = \frac{1}{(b-x)^2} \int_x^b f(s) ds + \frac{1}{(x-a)^2} \int_a^x f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)}$$

and

$$\begin{aligned}
 \frac{F(b) - F(a)}{b-a} &= \frac{1}{b-a} \left\{ f(b) - \frac{1}{b-a} \int_a^b f(s) ds - \left[\frac{1}{b-a} \int_a^b f(s) ds - f(a) \right] \right\} \\
 &= \frac{1}{b-a} \left\{ f(a) + f(b) - \frac{2}{b-a} \int_a^b f(s) ds \right\}
 \end{aligned}$$

therefore, we have

$$\begin{aligned}
 (2.3) \quad & F'(x) = \frac{1}{(b-x)^2} \int_x^b f(s) ds + \frac{1}{(x-a)^2} \int_a^x f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)} \\
 &= \frac{2}{b-a} \left[\frac{f(b) + f(a)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \right].
 \end{aligned}$$

Now, for $x \in (a, b)$, we set

$$M := \max \{x-a, b-x\} = \left[\frac{b-a}{2} + \left| x - \frac{a+b}{2} \right| \right],$$

and

$$m := \min \{x - a, b - x\} = \left[\frac{b-a}{2} - \left| x - \frac{a+b}{2} \right| \right],$$

so that, since f is positive we have

$$\frac{1}{M^2} \int_a^x f(s) ds \leq \frac{1}{(x-a)^2} \int_a^x f(s) ds \leq \frac{1}{m^2} \int_a^x f(s) ds$$

and

$$\frac{1}{M^2} \int_x^b f(s) ds \leq \frac{1}{(b-x)^2} \int_x^b f(s) ds \leq \frac{1}{m^2} \int_x^b f(s) ds$$

By adding the above two inequalities we get

(2.4)

$$\frac{1}{M^2} \int_a^b f(s) ds \leq \frac{1}{(b-x)^2} \int_x^b f(s) ds + \frac{1}{(x-a)^2} \int_a^x f(s) ds \leq \frac{1}{m^2} \int_a^b f(s) ds$$

and by (2.3) and (2.4) we get

$$\begin{aligned} & \frac{1}{M^2} \int_a^b f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)} \\ & \leq \frac{2}{b-a} \left[\frac{f(b)+f(a)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \right] \\ & = \frac{1}{(b-x)^2} \int_c^b f(s) ds + \frac{1}{(x-a)^2} \int_a^x f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)} \\ & \leq \frac{1}{m^2} \int_a^b f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)}. \end{aligned}$$

Hence, by multiplying the above inequality by the quantity $\frac{b-a}{2}$ we get

$$\begin{aligned} & \frac{b-a}{2} \left[\frac{1}{M^2} \int_a^b f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)} \right] \\ & \leq \frac{f(b)+f(a)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \\ & \leq \frac{b-a}{2} \left[\frac{1}{m^2} \int_a^b f(s) ds - \frac{(b-a)f(x)}{(x-a)(b-x)} \right]. \end{aligned}$$

Rearranging the terms we may write,

$$\begin{aligned} & \frac{(b-a)^2}{2M^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{M^2}{(x-a)(b-x)} f(x) \right] \\ & \leq \frac{f(b)+f(a)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \\ & \leq \frac{(b-a)^2}{2m^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{m^2}{(x-a)(b-x)} f(x) \right], \end{aligned}$$

for some $x \in (a, b)$; which proves the inequality (2.1). \square

Here, its convenient to note that

$$\begin{aligned}
 0 \leq \Delta &:= \text{The right-hand side of (2.1)} - \text{The left-hand side of (2.1)} \\
 &= \frac{(b-a)^2}{2m^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{m^2}{(x-a)(b-x)} f(x) \right] \\
 &\quad - \frac{(b-a)^2}{2M^2} \left[\frac{1}{b-a} \int_a^b f(s) ds - \frac{M^2}{(x-a)(b-x)} f(x) \right] \\
 (2.5) \quad &= \left(\frac{M^2 - m^2}{2m^2 M^2} \right) (b-a) \int_a^b f(s) ds,
 \end{aligned}$$

thus, it is clear that $\left(\frac{M^2 - m^2}{2m^2 M^2} \right) \geq 0$ and so that the difference $\Delta \geq 0$ iff $f(t) \geq 0$ $\forall t \in [a, b]$.

Finally, we note that another interesting form of the inequality (2.1) may be deduced by rewriting the terms of (2.1), to get:

$$\begin{aligned}
 (2.6) \quad &\frac{2M^2(b-a)}{(2M^2 + (b-a)^2)} \left[\frac{(b-a)^2}{2(x-a)(b-x)} f(x) + \frac{f(a) + f(b)}{2} \right] \\
 &\geq \int_a^b f(s) ds \\
 &\geq \frac{2m^2(b-a)}{(2m^2 + (b-a)^2)} \left[\frac{(b-a)^2}{2(x-a)(b-x)} f(x) + \frac{f(a) + f(b)}{2} \right],
 \end{aligned}$$

and so that, we have

$$\begin{aligned}
 (2.7) \quad 0 &\leq \int_a^b f(s) ds - \frac{2m^2(b-a)}{(2m^2 + (b-a)^2)} \Psi_f(a, b; x) \\
 &\leq \left[\frac{2M^2(b-a)}{(2M^2 + (b-a)^2)} - \frac{2m^2(b-a)}{(2m^2 + (b-a)^2)} \right] \Psi_f(a, b; x)
 \end{aligned}$$

where,

$$\Psi_f(a, b; x) := \frac{(b-a)^2}{2(x-a)(b-x)} f(x) + \frac{f(a) + f(b)}{2}$$

for some $x \in (a, b)$.

In an interesting particular case, let \mathcal{F} be the set of all functions $f : I \subset \mathbb{R} \rightarrow \mathbb{R}_+$ that satisfy the assumptions of Theorem 1 such that the required $x \in (a, b)$ is $x = \frac{a+b}{2}$, (in this case we have $M = m = \frac{b-a}{2}$) thus from (2.7) every such f satisfies that

$$(2.8) \quad \int_a^b f(s) ds = \frac{1}{3} (b-a) \left[2f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right],$$

where,

$$\Psi_f(a, b; \frac{a+b}{2}) = 2f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2}.$$

On the other hand, it is well-known that the error term in Simpson's quadrature rule (2.9) involves a fourth derivatives, however using the above observation and for every $f \in \mathcal{F}$; $\int_a^b f(s) ds$ can be evaluated 'exactly' using the Simpson formula (2.8) with no errors, and without going through its higher derivatives which may not exists or hard to find; as in the classical result.

$$(2.9) \quad \int_a^b f(x) dx = \frac{b-a}{3} \left[\frac{f(a)+f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] + \frac{(b-a)^5}{90} \|f^{(4)}\|_{\infty}.$$

3. APPLICATIONS TO MEANS

A function $M : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$, is called a Mean function if it has the following properties:

- (1) Homogeneity: $M(ax, ay) = aM(x, y)$, for all $a > 0$,
- (2) Symmetry : $M(x, y) = M(y, x)$,
- (3) Reflexivity : $M(x, x) = x$,
- (4) Monotonicity: If $x \leq x'$ and $y \leq y'$, then $M(x, y) \leq M(x', y')$,
- (5) Internality: $\min\{x, y\} \leq M(x, y) \leq \max\{x, y\}$.

We shall consider the means for arbitrary positive real numbers α, β ($\alpha \neq \beta$), see [6]–[7]. We take

- (1) The arithmetic mean :

$$A := A(\alpha, \beta) = \frac{\alpha + \beta}{2}, \quad \alpha, \beta \in \mathbb{R}_+.$$

- (2) The geometric mean :

$$G := G(\alpha, \beta) = \sqrt{\alpha\beta}, \quad \alpha, \beta \in \mathbb{R}_+$$

- (3) The harmonic mean :

$$H := H(\alpha, \beta) = \frac{2}{\frac{1}{\alpha} + \frac{1}{\beta}}, \quad \alpha, \beta \in \mathbb{R}_+ - \{0\}.$$

- (4) The power mean :

$$M_r(\alpha, \beta) = \left(\frac{\alpha^r + \beta^r}{2} \right)^{\frac{1}{r}}, \quad r \geq 1, \quad \alpha, \beta \in \mathbb{R}_+$$

- (5) The identric mean:

$$I(\alpha, \beta) = \begin{cases} \frac{1}{e} \left(\frac{\beta^\beta}{\alpha^\alpha} \right)^{\frac{1}{\beta-\alpha}}, & \alpha \neq \beta \\ \alpha, & \alpha = \beta \end{cases}, \quad \alpha, \beta > 0$$

- (6) The logarithmic mean :

$$L := L(\alpha, \beta) = \frac{\alpha - \beta}{\ln|\alpha| - \ln|\beta|}, \quad |\alpha| \neq |\beta|, \quad \alpha, \beta \neq 0, \quad \alpha, \beta \in \mathbb{R}_+.$$

- (7) The generalized log-mean:

$$L_p := L_p(\alpha, \beta) = \left[\frac{\beta^{p+1} - \alpha^{p+1}}{(p+1)(\beta - \alpha)} \right]^{\frac{1}{p}}, \quad p \in \mathbb{R} \setminus \{-1, 0\}, \quad \alpha, \beta > 0.$$

It is well known that L_p is monotonic nondecreasing over $p \in \mathbb{R}$, with $L_{-1} := L$ and $L_0 := I$. In particular, we have the following inequality $H \leq G \leq L \leq I \leq A$.

As a direct example on the inequality (2.1), consider $f : [a, b] \subset (0, \infty) \rightarrow (0, \infty)$ given by $f(s) = \frac{1}{s^2}$, $s \in [a, b]$, it is easy to see that $F(t) = \frac{b-a}{ab} \cdot \frac{1}{t^2}$, and so that the required $x \in (a, b)$ is $x = \sqrt{ab} := G(a, b)$. Applying (2.1), we get

$$(3.1) \quad \frac{(b-a)^2}{2M^2} \left[1 - \frac{M^2}{(G(a, b) - a)(b - G(a, b))} \right] \leq \frac{G^2(a, b)}{H(a^2, b^2)} - 1 \\ \leq \frac{(b-a)^2}{2m^2} \left[1 - \frac{m^2}{(G(a, b) - a)(b - G(a, b))} \right],$$

where, $M := \left[\frac{b-a}{2} + |G(a, b) - A(a, b)| \right]$, and $m := \left[\frac{b-a}{2} - |G(a, b) - A(a, b)| \right]$.

In general, the reader may check that it is not easy to find the value of x that satisfies the inequality (2.1). For example, we consider $f : [a, b] \subset (0, \infty) \rightarrow (0, \infty)$ given by

(1-) $f(s) = \frac{1}{s}$, $s \in [a, b] \subset (0, \infty)$, so that $F(t) = \ln \left(\frac{(b/t)^{1/(b-t)}}{(t/a)^{1/(t-a)}} \right)$. Applying (2.1), we get

$$(3.2) \quad \frac{(b-a)^2}{2M^2} \left[\frac{1}{L(a, b)} - \frac{M^2}{(x-a)(b-x)} \frac{1}{x} \right] \leq \frac{1}{H(a, b)} - \frac{1}{L(a, b)} \\ \leq \frac{(b-a)^2}{2m^2} \left[\frac{1}{L(a, b)} - \frac{m^2}{(x-a)(b-x)} \frac{1}{x} \right]$$

where $x \in (a, b)$ satisfying the equation (2.2), and m, M are defined above.

(2-) $f(s) = \ln(s)$, $s \in [a, b] \subset (0, \infty)$, so that

$$F(t) = \ln \left(\frac{(b^b/t^t)^{1/(b-t)}}{(t^t/a^a)^{1/(t-a)}} \right) = \ln \left(\frac{I(t, b)}{I(a, t)} \right), \quad t \in [a, b]$$

where $I(\cdot, \cdot)$ is the identric mean. Now, applying (2.1), we get

$$(3.3) \quad \frac{(b-a)^2}{2M^2} \left[\ln I(a, b) - \frac{M^2 \ln x}{(x-a)(b-x)} \right] \leq \ln G(a, b) - \ln I(a, b) \\ \leq \frac{(b-a)^2}{2m^2} \left[\ln I(a, b) - \frac{m^2 \ln x}{(x-a)(b-x)} \right]$$

where $x \in (a, b)$ satisfying the equation (2.2), and m, M are defined above.

(3-) $f(s) = s^p$, $s \in [a, b] \subset (0, \infty)$ and $p \in \mathbb{R} \setminus \{-1, 0\}$, so that

$$F(t) = \frac{b^{p+1} - t^{p+1}}{(b-t)(p+1)} - \frac{t^{p+1} - a^{p+1}}{(t-a)(p+1)} = L_p^p(t, b) - L_p^p(a, t), \quad t \in [a, b]$$

where $L_p^p(\cdot, \cdot)$ is the generalized logarithmic mean. Applying (2.1), we get

$$(3.4) \quad \frac{(b-a)^2}{2M^2} \left[L_p^p(a, b) - \frac{M^2 x^p}{(x-a)(b-x)} \right] \leq M_p^p(a, b) - L_p^p(a, b) \\ \leq \frac{(b-a)^2}{2m^2} \left[L_p^p(a, b) - \frac{m^2 x^p}{(x-a)(b-x)} \right]$$

where $x \in (a, b)$ satisfying the equation (2.2), and m, M are defined above.

REFERENCES

- [1] M.W. Alomari, A companion of the generalized trapezoid inequality and applications, *Journal of Math. Appl.*, 36 (2013), 5–15.
- [2] M.W. Alomari, New sharp inequalities of Ostrowski and generalized trapezoid type for the Riemann–Stieltjes integrals and applications, *Ukrainian Mathematical Journal*, 65 (7) (2013), 995–1018.
- [3] M.W. Alomari, M. Darus and U.S. Kirmaci, Some inequalities of Hermite-Hadamard type for s -convex functions, *Acta Mathematica Scientia*, 31 B(4) (2011) : 1643–1652.
- [4] M.W. Alomari, M. Darus and U. Kirmaci, Refinements of Hadamard-type inequalities for quasi-convex functions with applications to trapezoidal formula and to special means, *Comp. Math. Appl.*, 59 (2010), 225–232.
- [5] M. Alomari and M. Darus, On the Hadamard’s inequality for log-convex functions on the coordinates, *J. Ineq. Appl.*, 2009, Article ID 283147, 13 pages, doi:10.1155/2009/283147.
- [6] P. S. Bullen, D. S. Mitrinović and M. Vasić, Means and Their Inequalities, Dordrecht: Kluwer Academic, 1988.
- [7] P. S. Bullen, Handbook of Means and Their Inequalities, Dordrecht: Kluwer Academic, 2003.
- [8] S.S. Dragomir, Two mappings in connection to Hadamard’s inequalities, *J. Math. Anal. Appl.*, 167 (1992) 49–56.
- [9] S.S. Dragomir and R.P. Agarwal, Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula, *Appl. Math. Lett.*, 11 (1998) 91–95.
- [10] S. S. Dragomir and C. E. M. Pearce, “Selected Topics on Hermite-Hadamard Inequalities and Applications,” RGMIA Monographs, Victoria University, 2000, [http://www.staff.vu.edu.au/RGMIA/monographs/hermite hadamard.html](http://www.staff.vu.edu.au/RGMIA/monographs/hermite%20hadamard.html).
- [11] S.S. Dragomir, Y.J. Cho and S.S. Kim, Inequalities of Hadamard’s type for Lipschitzian mappings and their applications, *J. Math. Anal. Appl.*, 245 (2000), 489–501.
- [12] D.A. Ion, Some estimates on the Hermite-Hadamard inequality through quasi-convex functions, *Annals of University of Craiova, Math. Comp. Sci. Ser.*, 34 (2007), 82–87.
- [13] U.S. Kirmaci, Inequalities for differentiable mappings and applications to special means of real numbers to midpoint formula, *Appl. Math. Comp.*, 147 (2004), 137–146.
- [14] U.S. Kirmaci and M.E. Özdemir, On some inequalities for differentiable mappings and applications to special means of real numbers and to midpoint formula, *Appl. Math. Comp.*, 153 (2004), 361–368.
- [15] M.E. Özdemir, A theorem on mappings with bounded derivatives with applications to quadrature rules and means, *Appl. Math. Comp.*, 138 (2003), 425–434.
- [16] C.E.M. Pearce and J. Pečarić, Inequalities for differentiable mappings with application to special means and quadrature formula, *Appl. Math. Lett.*, 13 (2000) 51–55.
- [17] G.S. Yang, D.Y. Hwang and K.L. Tseng, Some inequalities for differentiable convex and concave mappings, *Comp. Math. Appl.*, 47 (2004), 207–216.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND INFORMATION TECHNOLOGY, IRBID NATIONAL UNIVERSITY, 21110 IRBID, JORDAN
E-mail address: mwomath@gmail.com